

Surface effects: convection dynamics & nonadiabaticity

Günter Houdek

Magnus Aarslev


Regner Trampedach

Jørgen Christensen-Dalsgaard


Surface effects: current models for adiabatic frequency corrections

Solar observations – **adiabatic** calculations

Christensen-Dalsgaard et al. (1996)




Rosenthal et al. (1995)



Surface effects: current models for adiabatic frequency corrections

Solar observations – **adiabatic** calculations




$$\frac{\partial r}{\partial m} = \frac{1}{4\pi r^2 \rho} \quad ; \quad \frac{\partial p}{\partial m} = -\frac{1}{4\pi r^2} \left(g + \frac{\partial r^2}{\partial t^2} \right)$$

$p = p_g + p_t$

Surface effects: current models for adiabatic frequency corrections

Solar observations – **adiabatic** calculations




$$\frac{\partial r}{\partial m} = \frac{1}{4\pi r^2 \rho} \quad ; \quad \frac{\partial p}{\partial m} = -\frac{1}{4\pi r^2} \left(g + \frac{\partial r^2}{\partial t^2} \right)$$

$p = p_g + p_t$

$$\frac{d}{dm} \left(\frac{\delta r}{r} \right) = -\frac{1}{4\pi r^3 \rho} \left(3 \frac{\delta r}{r} + \frac{\delta \rho}{\rho} \right)$$

Surface effects: current models for adiabatic frequency corrections

Solar observations – **adiabatic** calculations



$$\frac{\partial r}{\partial m} = \frac{1}{4\pi r^2 \rho} \quad ; \quad \frac{\partial \overset{p = p_g + p_t}{p}}{\partial m} = -\frac{1}{4\pi r^2} \left(g + \frac{\partial r^2}{\partial t^2} \right)$$


$$\frac{d}{dm} \left(\frac{\delta r}{r} \right) = -\frac{1}{4\pi r^3 \rho} \left(3 \frac{\delta r}{r} + \frac{\delta \rho}{\rho} \right)$$

Nonadiabatic EOS:

$$\frac{\delta \rho}{\rho} = \alpha \underbrace{\frac{\overset{p}{p_g}}{\left(\frac{\delta p}{p} - \frac{\delta p_t}{p} \right)}}_{\delta p_g / p_g} - \hat{\delta} \frac{\delta T}{T}$$

Surface effects: current models for adiabatic frequency corrections

Solar observations – **adiabatic** calculations



$$\frac{\partial r}{\partial m} = \frac{1}{4\pi r^2 \rho} \quad ; \quad \frac{\partial \overset{p = p_g + p_t}{p}}{\partial m} = -\frac{1}{4\pi r^2} \left(g + \frac{\partial r^2}{\partial t^2} \right)$$


$$\frac{d}{dm} \left(\frac{\delta r}{r} \right) = -\frac{1}{4\pi r^3 \rho} \left(3 \frac{\delta r}{r} + \frac{\delta \rho}{\rho} \right)$$

Nonadiabatic EOS: $\frac{\delta \rho}{\rho} = \alpha \underbrace{\frac{\overset{p}{p_g} \left(\frac{\delta p}{p} - \frac{\delta p_t}{p} \right)}_{\delta p_g / p_g} - \hat{\delta} \frac{\delta T}{T}$

Adiabatic EOS: $\frac{\delta \rho}{\rho} = \frac{1}{\gamma_1} \frac{\delta p_g}{p_g} = \frac{1}{\gamma_1} \frac{\overset{p}{p_g} \left(\frac{\delta p}{p} - \frac{\delta p_t}{p} \right)}{\overset{p}{p_g}}$

Surface effects: current models for adiabatic frequency corrections

Solar observations – **adiabatic** calculations



$$\frac{\partial r}{\partial m} = \frac{1}{4\pi r^2 \rho} \quad ; \quad \frac{\partial \overset{p = p_g + p_t}{p}}{\partial m} = -\frac{1}{4\pi r^2} \left(g + \frac{\partial r^2}{\partial t^2} \right)$$


$$\frac{d}{dm} \left(\frac{\delta r}{r} \right) = -\frac{1}{4\pi r^3 \rho} \left(3 \frac{\delta r}{r} + \frac{\delta \rho}{\rho} \right)$$

Nonadiabatic EOS: $\frac{\delta \rho}{\rho} = \alpha \underbrace{\frac{\overset{p}{p_g} \left(\frac{\delta p}{p} - \frac{\delta p_t}{p} \right)}_{\delta p_g / p_g} - \hat{\delta} \frac{\delta T}{T}$

Adiabatic EOS: $\frac{\delta \rho}{\rho} = \frac{1}{\gamma_1} \frac{\delta p_g}{p_g} = \frac{1}{\gamma_1} \frac{\overset{p}{p_g} \left(\frac{\delta p}{p} - \cancel{\frac{\delta p_t}{p}} \right)}{\overset{p}{p_g}}$

Surface effects: current models for adiabatic frequency corrections


Solar observations – **adiabatic** calculations



→ δp_t contributes predominantly to **imaginary part** of the (complex) pulsation frequency

Surface effects: current models for adiabatic frequency corrections

Solar observations – **adiabatic** calculations




$$\frac{\partial r}{\partial m} = \frac{1}{4\pi r^2 \rho} \quad ; \quad \frac{\partial \overset{p = p_g + p_t}{p}}{\partial m} = -\frac{1}{4\pi r^2} \left(g + \frac{\partial r^2}{\partial t^2} \right)$$

$$\frac{d}{dm} \left(\frac{\delta r}{r} \right) = -\frac{1}{4\pi r^3 \rho} \left(3 \frac{\delta r}{r} + \frac{\delta \rho}{\rho} \right)$$

Adiabatic EOS: $\frac{\delta \rho}{\rho} = \frac{1}{\gamma_1} \frac{\delta p_g}{p_g} = \frac{1}{\gamma_1} \frac{p}{p_g} \left(\frac{\delta p}{p} - \frac{\delta p_t}{p} \right)$

Surface effects: current models for adiabatic frequency corrections

Solar observations – **adiabatic** calculations



$$\frac{\partial r}{\partial m} = \frac{1}{4\pi r^2 \rho} \quad ; \quad \frac{\partial p}{\partial m} = -\frac{1}{4\pi r^2} \left(g + \frac{\partial r^2}{\partial t^2} \right)$$

$p = p_g + p_t$

$$\frac{d}{dm} \left(\frac{\delta r}{r} \right) = -\frac{1}{4\pi r^3 \rho} \left(3 \frac{\delta r}{r} + \frac{\delta \rho}{\rho} \right)$$


Adiabatic EOS: $\frac{\delta \rho}{\rho} = \frac{1}{\gamma_1} \frac{\delta p_g}{p_g} = \frac{1}{\gamma_1} \frac{p}{p_g} \left(\frac{\delta p}{p} - \frac{\delta p_t}{p} \right)$

Adiabatic EOS: $\frac{\delta \rho}{\rho} = \frac{1}{\gamma_1} \frac{\delta p_g}{p_g} \simeq \frac{1}{\gamma_1} \frac{p}{p_g} \frac{\delta p}{p}$

$\frac{1}{\gamma_1} \frac{p}{p_g}$

Surface effects: current models for adiabatic frequency corrections

Solar observations – **adiabatic** calculations



$$\frac{\partial r}{\partial m} = \frac{1}{4\pi r^2 \rho} \quad ; \quad \frac{\partial \overset{p = p_g + p_t}{p}}{\partial m} = -\frac{1}{4\pi r^2} \left(g + \frac{\partial r^2}{\partial t^2} \right)$$

$$\frac{d}{dm} \left(\frac{\delta r}{r} \right) = -\frac{1}{4\pi r^3 \rho} \left(3 \frac{\delta r}{r} + \frac{\delta \rho}{\rho} \right)$$

Adiabatic EOS: $\frac{\delta \rho}{\rho} = \frac{1}{\gamma_1} \frac{\delta p_g}{p_g} = \frac{1}{\gamma_1} \frac{\overset{p}{p_g}}{\overset{p_g}{p}} \left(\frac{\delta p}{p} - \cancel{\frac{\delta p_t}{p}} \right)$


$\frac{1}{\gamma_1}$

Adiabatic EOS: $\frac{\delta \rho}{\rho} = \frac{1}{\gamma_1} \frac{\delta p_g}{p_g} \simeq \underbrace{\frac{1}{\gamma_1} \frac{\overset{p}{p_g}}{\overset{p_g}{p}}}_{\frac{1}{\gamma_1} \frac{\overset{p}{p_g}}{\overset{p_g}{p}}} \frac{\delta p}{p}$

$\frac{1}{\gamma_1} \frac{\overset{p}{p_g}}$


1D models including turbulent pressure p_t

$$\frac{\partial}{\partial m}(p_g + p_t) = -\frac{1}{4\pi r^2} \left(\frac{Gm}{r^2} + \frac{\partial^2 r}{\partial t^2} \right)$$




1D models including turbulent pressure p_t

$$\frac{\partial}{\partial m}(p_g + p_t) = -\frac{1}{4\pi r^2} \left(\frac{Gm}{r^2} + \frac{\partial^2 r}{\partial t^2} \right)$$




1D models with turbulent pressure p_t , δp_t , $\delta(L_r + L_c)$



weak nonlocality (GH 1997)

1D models with turbulent pressure p_t , δp_t , $\delta(L_r + L_c)$




weak nonlocality (GH 1997)

1D models with turbulent pressure p_t , δp_t , $\delta(L_r + L_c)$

Nonadiabaticity: $\delta(L_r + L_c)$

Convection dynamics: δp_t




weak nonlocality (GH 1997)

1D models with turbulent pressure p_t , δp_t , $\delta(L_r + L_c)$

Nonadiabaticity: $\delta(L_r + L_c)$

Convection dynamics: δp_t




weak nonlocality (GH 1997)

1D models with turbulent pressure p_t , δp_t , $\delta(L_r + L_c)$

Nonadiabaticity: $\delta(L_r + L_c)$

Convection dynamics: δp_t




Patched 1D models with turbulent pressure p_t , δp_t , $\delta(L_r + L_c)$


MDI observations — 1D.PM[p_t].adiabatic

“Patched Model”

M. Aarslev (2016)




Patched 1D models with turbulent pressure p_t , δp_t , $\delta(L_r + L_c)$




M. Aarslev (2016)

calibrated max p_t (strong nonlocality)

1D models including turbulent pressure p_t




Patched 1D models with turbulent pressure p_t , δp_t , $\delta(L_r + L_c)$



calibrated max p_t (strong nonlocality)


1D model with turbulent pressure p_t , δp_t , $\delta(L_r + L_c)$



calibrated max p_t (strong nonlocality)

1D model with turbulent pressure p_t , δp_t , $\delta(L_r + L_c)$

MDI observations — 1D[$p_t, \delta p_t, \delta F_c$].na





$$\frac{\delta \rho}{\rho} \simeq \frac{1}{\gamma_1} \frac{p}{p_g} \frac{\delta p}{p}$$



$$\frac{\delta \rho}{\rho} \simeq \frac{1}{\gamma_1} \frac{\delta p}{p}$$

calibrated max p_t (strong nonlocality)

1D model with turbulent pressure p_t , δp_t , $\delta(L_r + L_c)$



1D model with turbulent pressure p_t , δp_t , $\delta(L_r + L_c)$



Points still to consider

- effect of turbulent **velocity anisotropy**
- modal effects due to **advection** of the oscillations by **spatially varying** radial **flows**
Zhugzhda & Stix (1994, A&A)
Bhattacharya, Hanasoge, Antia (2015, ApJ)
- Calculate “full” nonadiabatic (with convection dynamics) directly from the patched 1D/<3D> models
- effects we haven’t thought about

Surface effect: convection dynamics and nonadiabaticity

Anisotropy factor:

$$\Phi = \frac{\langle u^2 \rangle + \langle v^2 \rangle + \langle w^2 \rangle}{\langle w^2 \rangle}$$

$\mathbf{u} = (u, v, w)$... turbulent velocity field

